
Outcomes for Dutch patients at Castle Craig Hospital



the 2012 evaluation for all Dutch patients admitted between 13.04.09 to 27.12.10

Independent analysis of outcome data
Christo Research Systems

31st October 2012

Castle Craig Hospital

Medical Director: Dr Margaret McCann MB, Bch, BAO.
Treatment Director: Dr Florian Kaplick Psychiatrist, PhD.
Address: Blyth Bridge, West Linton, Peeblesshire, Scotland EH46 7DH.
Tel: 01721 722763
E-mail: enquiries@castlecraig.co.uk

Christo Research Systems

Director: Dr George Christo DipHE, BSc, PhD, PsychD, CPsychol, AFBPsS.
Tel: 07754 786 379
E-mail: DrGeorgeChristo@tiscali.co.uk
Website: <http://users.breathemail.net/drgeorgechristo/>

Summary of findings

- One hundred and six drug and 95 alcohol dependent patients from the Netherlands entered Castle Craig Hospital from 13th April 2009 to 27th December 2010 and stayed in treatment for more than 1 day.
- All 201 patients were Dutch nationals from the Netherlands, 178 males and 23 females.
- Most patients were alcohol or cocaine addicts.
- The patients' average age was 38.0 years (*range* = 18 - 68). Females were generally older than the males, drinkers tended to be older and cocaine users tended to be younger.
- One hundred and seventeen of the patients had been formally diagnosed with another mental health condition in addition to their substance misuse.
- Most patients (135) were self-referred, 42 were referred by their General practitioner, 17 from mental health or addictions services, and 7 from psychiatrists.
- Those referred from mental health or addictions services were less likely to complete treatment. Those referred by their GP were far more likely to complete.
- Patients intake levels of dysfunction were similar to drinkers and drug users attending outpatient services (based on the CISS comparison scores for these groups, see Appendix). Patients' greatest problems were with drug or alcohol use, lack of support, lack of occupation and psychological problems.
- The patients' average wait between referral to treatment entry was 8.7 weeks.
- 74% of patients completed treatment, 18% of patients prematurely self-discharged against medical advice, and 8% of patients were prematurely discharged for rule violations.
- Only 26 of the 201 patients (12.9%) stayed less than 6 weeks which is the usual duration of primary treatment.
- The average treatment duration for patients completing treatment was 19 weeks, and the average treatment duration for prematurely discharged patients was 8 weeks.
- Premature treatment discharge was more likely among patients also diagnosed with a personality disorder.
- Almost two thirds of patients completing treatment did well, whereas only one third taking premature discharges (for whatever reason) had a good outcome.
- Follow-ups were successfully completed on 156 of the 201 patients (78% response rate). Thirty six patients could not be contacted, 5 patients refused to answer the follow-up questions, 2 patients had died, 1 patient was in another addictions clinic, and 1 patient was unable to answer questions because of a psychiatric condition.

- The average follow-up period was 62 weeks after discharge.
- 82% of followed-up patients improved, 13% remained the same and 5% got worse.
- Good outcome was predicted by
 - Completion of treatment
 - Longer treatment duration
 - Fewer psychological problems
 - Fewer compliance problems
 - Fewer working relationship problems
- The average intake CISS score of the 156 followed-up patients was 9.7 and the average follow-up CISS score was 4.4 indicating a highly significant improvement in general functioning.
- Reductions in drug / alcohol use at follow-up were accompanied by improvements in all other CISS domains.
- Even those who were not totally abstinent at follow-up appeared to have benefited from their experience in treatment, probably by gaining a period of respite during which to recover from the consequences of their excessive drinking or drug use.
- Post treatment '12 step' meeting attendance was associated with good outcomes.
- The following success rates are conservatively based by including all 201 patients on the assumption that the 45 patients not followed-up (response rate = 78%) showed no improvement or otherwise had poor outcomes.
 - Being totally abstinent from all drugs or alcohol at follow-up
49.3% ($n = 99$)
 - Achieving low problem severity at follow-up (CISS < 6, see appendix)
54.2% ($n = 109$)
 - Showing any reduction in measured levels of dysfunction
63.7% ($n = 128$)

However the success rates might be a little higher because some of the missing patients would not have been contactable due to having recovered and being in full time occupation.

- The following success rates are thus more liberally based by excluding the 43 patients not responding to follow-up. Although 1 patient in another addictions clinic, and 1 patient unable to answer questions because of a psychiatric condition, remained included as poor outcomes (*new sample size* = 158).
 - Being totally abstinent from all drugs or alcohol at follow-up
62.6% ($n = 99$)
 - Achieving low problem severity at follow-up (CISS < 6, see appendix)
69.0% ($n = 109$)
 - Showing any reduction in measured levels of dysfunction
81.0% ($n = 128$)

Treatment Overview

General approach

Castle Craig Hospital provides an abstinence oriented residential treatment for alcohol or drug dependent individuals. It uses an established treatment model developed in the US around 1950 and first imported to the UK in 1974 (Cook, 1988a). Outcomes generated by this approach are very good (Cook, 1988b) and have been shown to be at least equal to and in some cases better than other commonly used treatments for substance misuse (Project MATCH, 1997; Ouimette et al, 1997; Longabaugh et al, 1998).

It is an intensive psychologically oriented approach consisting of regular group work, one to one counselling, lectures and written assignments. Many similar therapeutic communities are well established throughout the UK and their programme facilitates engagement with the independent free after care resource provided by Alcoholics Anonymous and Narcotics Anonymous (AA & NA) groups. Regular attendance of AA and NA has been shown to be associated with reduced drug or alcohol use (Emrick, 1987; McLatchie & Lomp, 1988; Alford et al, 1991; Christo & Franey, 1995; Gossop et al, 2003), improved psychological health (Christo & Sutton, 1994; DeSoto et al, 1989; DeSoto et al, 1985; McCown, 1989; McCown, 1990), and with improved physical health (Mann et al, 1991).

Services offered

Castle Craig Hospital offers detoxification from alcohol, tranquillisers, or opiates. Patients are encouraged to engage with all aspects of the programme during detoxification because it serves as a useful distraction from withdrawal symptoms and assists in their orientation. Patients are also assessed to identify specific medical (e.g. liver dysfunction), psychological (e.g. cognitive deficits, anxiety, abuse or traumatic events), or psychiatric (e.g. dual diagnosis, suicide risk, epilepsy) problems that may need to be addressed in their individual care plans.

The first phase of treatment includes comprehensive assessment, detoxification, stabilisation, treatment planning and intensive individual and group psychotherapy. The duration of this phase is about six weeks. Therapy staff employ a full range of psychotherapeutic approaches depending upon their training and interests (e.g. Twelve Step Facilitation, Rational Emotive Behavioural Therapy, Cognitive Behavioural Therapy, Reality Therapy, and Transactional Analysis).

This phase continues with a second phase of short to medium term treatment with a further duration of from 8 to 24 weeks. This additional period of treatment is especially suitable for those patients with co morbid psychiatric disorders such as personality disorder, depressive disorders, phobias, generalised anxiety, poor support networks or social functioning. Addiction complicated by a psychiatric diagnosis is generally associated with worse outcomes. Such patients have more complex needs and need longer in treatment. The treatment in the second phase remains intensive, continuing to address those factors which have an immediate bearing on relapse. As well as using the same elements found in the primary phase of treatment, this second phase of care also assists patients to re-integrate with society by focusing on practical

issues of occupation, and family problems.

Castle Craig Hospital can offer aftercare group therapy held in a number of locations in Scotland and in the Netherlands at Amsterdam and Den Haag. All clients are encouraged to attend one or more of these aftercare sessions weekly for a period of up to two years after completing their residential treatment.

Aims of treatment

- Detoxification and stabilisation, abstaining from alcohol and other drugs.
- Creating a therapeutic bond to facilitate engagement with support from staff, peers, and AA or NA.
- Separating from people, places and things that promote substance use and establishing a new social network that supports recovery.
- Identifying recurrent problems, resolving painful / traumatic memories.
- Stopping compulsive self-defeating behaviours that suppress awareness of painful feelings and irrational thoughts.
- Relapse warning sign identification and management strategies. Identifying past causes of lapse and appropriate future coping strategies.
- Learning how to manage feelings and emotions responsibly without resorting to compulsive behaviour or the use of chemicals.
- Identifying and changing dysfunctional core beliefs (about self, others, and the world) that promote the use of irrational thinking and create painful feelings and self-defeating behaviours.
- Learning to change maladaptive behaviour patterns developed during childhood in dysfunctional families of origin.
- Increasing self-esteem by feeling worthwhile to self and helping others, promoting engagement with society, dealing with practical problems and establishing meaningful occupation.

Outcome measures & methods

How outcome was measured

Outcome was measured by the Christo Inventory for Substance-misuse Services (CISS) which is a standardised, validated tool (Christo, Spurrell & Alcorn, 2000, Christo, 2000a) now commonly used in Scotland (Effective Interventions Unit, 2001), England & Wales (Audit Commission, 2002; Christo, 1999a,b,c; Christo, 2000b,c,d,e,f, Christo, 2001), and abroad (Christo & Da Silva, 2002). The CISS is a single page outcome evaluation tool completed by drug / alcohol service workers either from direct client interviews or from personal experience of their client supplemented by existing assessment notes. Its purpose is to elicit workers' impressions of their clients in a quick, quantitative, standardised and reliable way. The 0 to 20 scale consists of 10 items reflecting clients' problems with:

Social functioning	Criminal involvement
General health	Drug / alcohol use
Sexual / injecting risk behaviour	Ongoing support
Psychological functioning	Compliance
Occupation	Working relationships

These outcome areas are scored on a three point scale of problem severity (0 = none, 1 = moderate, 2 = severe), each point is illustrated with relevant examples for guidance. Thus, a CISS score of 0 would indicate no problems and a score of 20 would indicate severe problems in all outcome areas.

Evaluation procedure

CISS is incorporated as a regular part of Castle Craig Hospital's intake and follow-up procedures. Baseline CISS forms were completed by staff from information gathered at the first assessment. They were then completed again during follow-up interviews on average about 62 weeks after discharge from treatment. A table of relevant dates, CISS information and other data for all Dutch patients was delivered to Christo Research Systems for analysis.

Sample

The sample comprised of all patients from the Netherlands who entered treatment between 13th April 2009 to 27th December 2010 and stayed in treatment for more than 1 day. Two hundred and one patients met these criteria, attempts were made to follow up all of them and 156 patients (77.6%) were successfully contacted in order to obtain the detailed information presented below. This evaluation thus details the outcomes for the 156 patients (137 males, 19 females) who were followed-up.

Findings regarding all 201 patients

Statistical information

- ***n*** indicates the number of individuals contributing to each statistical sample.
- ***m*** indicates a mean value, all averages in this report are means.
- ***sd*** indicates a standard deviation, thus giving an idea of the spread of scores around the mean. (In a normal distribution, 68% of all data points lie plus or minus one sd about the mean.)
- ***range*** indicates the total range of values within a measured variable (minimum - maximum).
- ***t*, *f*, χ^2** and ***U*** are statistical tests to show if groups are significantly different from each other.
- ***p*** indicates the level of significance of a statistical test, the smaller the better.

Gender

178 males	88.6 %
23 females	11.4 %

Nationality and ethnic origins: All 201 patients were Dutch nationals from the Netherlands.

Drugs of choice

95 patients using alcohol	47.3%	Alcohol 47.3%
69 patients using cocaine	34.3%	Drugs 52.7%.
18 patients using cannabis	9.0 %	.
12 patients using opiate/oids	6.0 %	.
5 patients using amphetamines	2.5 %	.
2 patients using sedatives	1.0 %	.

Secondary drugs of choice

Among the 95 drinkers, 62 of them had a secondary drug recorded as follows:

20 also using polysubstances

13 also using nicotine (although the majority of patients also smoke nicotine)

10 also using cocaine

11 also using cannabis

6 also using sedatives

2 also using opiate/oids

Among the 106 drug users, 92 of them had a secondary drug recorded as follows:

36 also using polysubstances

25 also using alcohol

12 also using cannabis

10 also using nicotine (although the majority of patients also smoke nicotine)

4 also using cocaine

3 also using sedatives

2 also using opiate/oids

The above picture of polysubstance use illustrates why interventions targeting single drug types might sometimes prove ineffective. Castle Craig's therapeutic target is total abstinence from all recreational drugs including alcohol (but not including nicotine and caffeine).

Table 1, Gender distribution by drug type

	Alcohol dependent	Cocaine dependent	Other	Total
Females	13 (56.5%)	5 (21.7%)	5 (21.7%)	23
Males	82 (46.1%)	64 (36.0%)	32 (18.0%)	178
Total	95	69	37	201

Table 1 above illustrates that there is no significant relationship between gender and drug of choice in this sample.

Table 2, Average Age by Drug type by Gender

The patients' average age was 38.0 years ($n = 201$, $sd = 10.8$, $range = 18 - 68$)

	Alcohol dependent	Cocaine dependent	Other drug	Total
Females	49.2 yrs ($n=13$)	46.3 yrs ($n=5$)	37.4 yrs ($n=5$)	46.0 yrs ($n=23$)
Males	42.3 yrs ($n=82$)	31.2 yrs ($n=64$)	34.7 yrs ($n=32$)	36.8 yrs ($n=178$)
Total	43.3 yrs ($n=95$)	32.3 yrs ($n=69$)	35.0 yrs ($n=37$)	38.0 yrs ($n=201$)

As illustrated in table 2 above, females were generally older than males ($t [199] = 3.9$, $p < .001$). Drinkers tended to be older and cocaine users tended to be younger ($f [2] = 28.4$, $p < .001$).

Table 3, Psychiatric Diagnoses by Gender, Drug type, & treatment completion

One hundred and seventeen of the patients had been formally diagnosed with another mental health condition in addition to their substance misuse.

Concurrent General Diagnosis	Detailed Concurrent Diagnoses	N	% female	% primary alcohol	% completed treatment
Nil		84	11%	50%	81%
Personality Disorders	Mostly Cluster B: Dramatic or Erratic Behaviours Borderline, narcissistic and antisocial	44	16%	57%	59%
Depressive Disorders	Mostly major depressive disorder, and some mood disorder and bipolar disorder	17	12%	47%	77%
Attention Deficit	Mostly attention deficit hyperactivity disorder	17	0%	18%	71%
Anxiety Disorders	Generalised anxiety, post traumatic stress syndrome, panic attacks, and social phobia	25	20%	44%	80%
Other	Cognitive disorder, eating disorder, adjustment disorder, autism, Rett's syndrome	7	0%	71%	71%
Gambling	Pathological gambling	4	0%	25%	75%
Psychosis	Psychotic disorders	3	0%	0%	33%
Total		201	11%	47%	74%

As illustrated in table 3 above; the overall proportion of female patients is 11% and a greater proportion of females than thus expected were found within the anxiety category. A lower proportion of females than expected were found within the attention deficit category. It also appeared that patients with personality disorder were more likely to leave treatment prematurely. However, there were too many categories with small cell counts to allow a meaningful statistical test.

Table 4, Psychiatric categories, treatment duration and treatment completion

Detailed mental health diagnoses were collapsed into 'other diagnoses' and 'personality disorder' categories for the purpose of analysis.

Referral Type	n	%	Mean treatment duration (weeks)	% completed treatment
Nil	84	41.8	16.9	81.0
Other diagnoses	73	36.3	16.3	74.0
Personality disorder	44	21.9	13.5	59.1
Total	201	100	15.9	73.6

Mean treatment duration was not significantly shorter for 'personality disordered' patients ($f [2] = 2.4, p = .09$). But they had the smallest percentage treatment completion rate ($\chi^2 [2] = 7.1, p = .03$).

Table 5, Referral sources

The greatest proportion of patients was self-referred followed by those referred by General Practitioners referred. However, all self-referred patients were instructed to get a referral letter from their GP, given that a GP referral is compulsory before treatment can begin.

Referral Source	<i>n</i>	%	Mean treatment duration (weeks)	% completed treatment	Mean age (years)	% female
Self-referred	135	67.2	14.8	70.4	37.2	9.6
General Practitioner (GP)	42	20.9	19.6	90.5	37.6	11.9
Mental health or addictions service	17	8.5	14.3	52.9	40.6	5.9
Psychiatrist	7	3.5	19.2	85.7	48.5	57.1
Total	201	100	15.9	73.6	38.0	11.4

Referral sources were not significantly related to drug type, or time taken to enter treatment.

Mean treatment duration was shortest for 'Mental health or addictions service' referrals and longest for referrals from General Practitioners ($f[3] = 4.1, p = .007$).


Percentage treatment completion was the least for 'Mental health or addictions service' referrals and the greatest for referrals from General Practitioners ($\chi^2 [3] = 11.2, p = .01$)

Psychiatrists appeared to have referred a greater proportion of females (likelihood ratio [3] = 9.6, $p = .02$). Although the numbers of females in this sample were relatively small anyway.

Psychiatrists' referrals also tended to be older ($f [3] = 2.8, p = .04$), and this may be related to the earlier finding that females in this sample tended to be older.

Time from referral to treatment entry (time to enter treatment)

Figure 1 below shows the patients' average time between referral and treatment entry was 8.7 weeks ($n = 201$, $sd = 5.5$, $range = 0.3 - 36.1$). Entry time was unrelated to gender, referral source, drug of choice, premature treatment discharge or eventual outcome.



The longer delays between referral to treatment entry were generally due to patients' issues such as:

- Ten patients needed a long detoxification in a clinic in the Netherlands before treatment
- Seven patients had problems with their insurance company that needed to be solved before they could enter treatment
- Five patients had a lengthy wait between assessment and admission because the treatment was started as outpatient.
- Two patients were ambivalent in their motivation to quit and it took them a long time to make up their mind
- Two patients had already planned holidays and took those before entering treatment
- Two patients were delayed because of ongoing legal issues
- Two patients wanted to delay their treatment entry
- Two patients had delays between assessment and admission for reasons unknown
- One patient was reluctant to get a referral letter from his GP
- One patient was unable to travel by airplane
- One patient wanted to wait for an appointment for a job interview
- One patient was delayed by work related issues
- One patient had to go to hospital after a drug overdose
- One patient was delayed because of financial problems
- One patient was delayed because of medical problems

Patients' problems at intake

The average intake CISS total score of the 201 patients was 9.9 (*sd* = 3.1, *range* 4 - 19) and there was no significant relationship between gender, drug of choice and average total score. This figure is indicative of an average level of dysfunction and suggests that these patients are similar in dysfunction to drinkers and drug users attending outpatient services (based on the CISS comparison scores for these groups, see Appendix). Patients' greatest problems were with drug or alcohol use, lack of support, lack of occupation and psychological problems.

For 106 drug dependent patients:

- 3.8 % of patients had low problem severity (CISS score 0 to 5)
- 75.5% of patients had average problem severity (CISS score 6 to 12)
- 20.8% of patients had high problem severity (CISS score 13 to 20)

For 95 alcohol dependent patients:

- 1.1% of patients had low problem severity (CISS score 0 to 4)
- 72.6% of patients had average problem severity (CISS score 5 to 11)
- 26.3% of patients had high problem severity (CISS score 12 to 20)

Figure 2, Baseline CISS item scores by drug type




Figure 2 compares the average CISS item scores (0 to 2 scale) as assessed at intake between the 106 drug and the 95 alcohol dependent patients. The taller the bars in the figure, the greater the average degree of problem within the relevant CISS domain. No differences emerged between drug or alcohol dependent groups.

No gender differences emerged among the intake CISS domains between drug or alcohol dependent groups.

Reasons for discharge from Castle Craig

- 148 Patients completed treatment 73.6 %
- 53 Patients left prematurely (incomplete treatment) 26.4 %
- of which...
 - 37 patients prematurely self-discharged against medical advice 18.4 %
 - 16 patients were prematurely discharged for rule violations 8.0 %

The average treatment duration for patients completing treatment was 18.6 weeks ($n = 148$, $sd = 7.5$), it was significantly longer than the prematurely discharged patients' average treatment duration of 8.4 weeks ($n = 53$, $sd = 6.4$), ($t [199] = 8.9$, $p < .001$).

- There was no relationship between intake CISS scores and premature discharge.
- There was no relationship between age and premature discharge.
- There was no relationship between gender and premature discharge.
- There was no relationship between drug of choice and premature discharge.
- There was no relationship between insurance company and premature discharge.
- There was a relationship between referral source and premature discharge (see page 10).
- There was a relationship between concurrent personality disorder diagnoses and premature treatment discharge (see page 9).

Dropout rate during the first six weeks

Only 26 of the 201 patients (12.9%) stayed less than 6 weeks which is the usual duration of primary treatment.

Outcome

Attempts were made between 22.02.11 and 23.02.12 to contact all 201 patients via telephone, e-mail and post. Those who could be contacted were interviewed using the CISS outcome measure. Follow-ups were successfully completed on 156 of the 201 patients (77.6%). Two patients died and of the remaining 43:

- 36 patients could not be contacted by telephone, e-mail or letter (they either moved, changed their telephone number or e-mail address and didn't leave a forwarding address or give their new number. Or they didn't want contact or they didn't have the time)
- 5 patients refused to answer the follow-up questions
- 1 patient was in another addictions clinic in South Africa
- 1 patient was unable to answer questions because of a psychiatric condition

In treatment outcome follow-up studies such as this one it is not unreasonable to assume that the majority of patients not accessible, will have relapsed.

Christo, Spurrell & Alcorn (2000) found a CISS cut-off score of 6 or less can be used to indicate "good outcome" for abstinence based treatment among drug users. This correctly identified 88% of outcomes where drug use was assessed only in month before follow-up, and 84% of outcomes where drug use was assessed over their entire six-month follow-up period. Alcoholics are generally expected to score one CISS point less than drug addicts.

So due to the large number of alcoholics in this sample, a conservative cut-off score of 5 or less was used to indicate "good outcome" for abstinence based treatment among the Castle Craig Patients.

The patients could thus be categorised under the following outcomes:

109 had a follow-up CISS score of 5 or less	54.2 %	Good outcome 54 %
47 had a follow-up CISS score of 6 or more	23.4 %	Poor outcome 46%
43 could not be contacted	21.4 %	.
2 patients died	1.0%	.

Table 6, Outcome by treatment completion, and treatment duration

Outcome	% Completed (<i>n</i>)	Mean weeks in treatment
Good (<i>n</i> = 109)	61.5% (91)	17.5 (<i>sd</i> = 8.1)
Poor (<i>n</i> = 92)	38.5% (57)	14.1 (<i>sd</i> = 8.7)
Significance	$\chi^2 [1] = 11.9, p = .001$	$t [199] = 2.9, p = .004$

Table 6 above illustrates that ‘good outcome’ is experienced by a greater proportion of those who completed their treatment; ‘good outcome’ is also associated with a longer stay in treatment.

Patients’ referral source, age, gender, drug of choice, mental health diagnosis, or time taken to enter treatment, were unrelated to outcome.

CISS scores at treatment entry and subsequent outcome

The mean CISS score for patients with a good outcome was 9.6 (*n* = 109, *sd* = 2.9), it was not significantly different from the poor outcome patients’ mean score of 10.3 (*n* = 92, *sd* = 3.2).

Figure 3, Baseline CISS item scores by outcome




Figure 3 compares the average CISS item scores (0 to 2 scale) as assessed at intake between the 109 good and the 92 poor outcome patients. The taller the bars in the figure, the greater the average degree of problem within the relevant CISS domain. ‘Good outcome’ patients tended to have fewer psychological problems ($U = 4306, p = .038$), fewer compliance problems ($U = 4108, p = .008$), and fewer working relationship problems ($U = 4241, p = .033$).

Table 7, Discharge status and outcome

	Completed treatment	Left Against Medical Advice	Discharged for rule violation	total
% good	61.5	35.1	31.3	109
% poor	38.5	64.9	68.8	92
Total number	148	37	16	201

The expected split in table 7 above should be 54.2% good : 45.8% poor for all discharge types. Notable departures from this indicate that almost two thirds (61.5%) of patients completing treatment do well, whereas only one third taking premature discharges (for whatever reason) have a good outcome. This finding is statistically significant ($\chi^2 [2] = 12.0, p = .003$). Future patients thinking of quitting treatment early should be made aware of this statistic.

Summary, predictors of planned discharge and good outcome

Premature discharge from treatment was associated with shorter treatment durations and poorer outcomes. But there were few predictors of premature discharge, only a tendency for personality disordered patients and those referred from mental health or addictions services to leave prematurely. However these factors did not predict eventual treatment outcome.

Eventual good outcome was predicted by

- Completion of treatment
- Longer treatment duration
- Fewer psychological problems
- Fewer compliance problems
- Fewer working relationship problems

However, it should be remembered that these are only statistical trends and many clients without the above qualities will have good outcomes in any case. For example, 35% of patients taking a premature discharge against medical advice went on to have a good outcome anyway.

Findings regarding 156 patients followed-up

Attempts were made between 22.02.11 and 23.02.12 to contact all 201 patients via telephone, e-mail and post. Those who could be contacted were interviewed using the CISS outcome measure. Follow-ups were successfully completed on 156 of the 201 patients (77.6%). Two patients died and of the remaining 43:

- 36 patients could not be contacted by telephone, e-mail or letter (they either moved, changed their telephone number or e-mail address and didn't leave a forwarding address or give their new number. Or they didn't want contact or they didn't have the time)
- 5 patients refused to answer the follow-up questions
- 1 patient was in another addictions clinic in South Africa
- 1 patient was unable to answer questions because of a psychiatric condition

Figure 4, follow-up periods




Figure 4 indicates the number of patients falling within each follow-up period. The majority of patients were followed-up after 50 weeks and the distribution is skewed to the longer follow-up periods on the right.

Standard follow-up times were hard to implement due to the long sampling period required to

capture all the Dutch patients' treatment entry dates.

- Treatment entry took place between 28.10.09 to 27.12.10
- Follow-up interviews took place between 22.02.11 and 23.02.12

The average follow-up period was...

- From treatment entry: 74.6 weeks ($n = 156$, $sd = 10.7$, $range = 53.9 - 107.9$)
- From treatment discharge: 57.9 weeks ($n = 156$, $sd = 5.9$, $range = 49.4 - 84.1$)

Changes in patient dysfunction at follow-up

Figure 5, reductions of patient dysfunction




Figure 5 illustrates the reductions in CISS total scores achieved by the 156 patients who were followed-up. The inter-rater reliability of the CISS (Christo et al., 2000) would indicate that a score fluctuation of plus or minus one point is attributable to variations of CISS interpretation between raters. As such, only changes of 2 or more points are recognised as 'genuine' and on that basis:

- 82.1% of patients improved
- 12.8% of patients remained the same
- 5.1% of patients got worse

Twenty four patients achieved reductions of 10 CISS points or more. Changes of this magnitude are not uncommon among those who achieve total abstinence but would likely be perceived by the patients and their significant others as nothing short of miraculous.

Figure 6, the process of change




Figure 6 displays how CISS total scores are distributed among the 156 patients. Dark bars indicate the score distributions at intake and the light bars indicate score distributions at follow-up.

The average intake CISS total score of the 156 patients was 9.7 ($sd = 3.0$, range 4 - 18)
The average follow-up CISS total score of the 156 patients was 4.4 ($sd = 3.3$, range 0 - 17)
A paired sample t-test indicates this reduction to be highly significant ($t [155] = 16.4$, $p < .001$)

Figure 7, Changes in individual CISS item scores




Figure 7 compares the average CISS item scores (0 to 2 scale) as assessed at intake and then again at follow-up. The taller the bars in the figure, the greater the average degree of problem within the relevant CISS domain. As can also be seen in figure 2, the greatest problems at intake were with drug or alcohol use, lack of support, lack of occupation, and psychological problems.

Ten Wilcoxon Signed Ranks statistical tests indicated that the reductions in all of the 10 CISS outcome domains were highly significant. Thus indicating that reductions in drug / alcohol use were generally accompanied by improvements in all other aspects of the patients' lives.

Table 8, Outcome and '12 step' meeting attendance reported at follow-up

Outcome	% Attenders (n)	% Non-attenders (n)
Good (n = 109)	83.5% (71)	59.4% (38)
Poor (n = 40)	16.5% (14)	40.6% (26)
Total n = 149	100% (85)	100% (64)

Table 8 above illustrates that '12 step' meeting attendance (e.g. Alcoholics Anonymous, Narcotics Anonymous, Cocaine Anonymous, etc.) data were available for 149 of the 156 followed-up patients. Within this sub-group, 83.5% of attenders and 59.4% of non-attenders were recorded as having a good treatment outcome. '12 step' meeting attendance is thus associated with good outcomes ($\chi^2 [1] = 10.8, p = .001$). Among the 85 attenders (mean attendance = 2.1 meetings per week, $sd = 1.5, range = 0.25$ to 7.0), there was no relationship between outcome and frequency of attendance (Spearman's rho [83] = $.03, p = .8$).

Lapses reported at follow-up

Lapse (back to drug / alcohol use) information was available for 151 of the 156 followed-up patients. Lapse frequency categories were described as follows:

1. 70 patients (46.4%) reported having no lapses since leaving treatment
2. 25 patients (16.6%) reported having 1 lapse since leaving treatment
3. 14 patients (9.3%) reported having 2 lapses since leaving treatment
4. 11 patients (7.3%) reported having 3 to 6 lapses since leaving treatment
5. 31 patients (20.5%) reported having 10 or more lapses since leaving treatment

Higher lapse frequency was predicted at baseline by:

- Being an alcohol user (*Spearman's rho* [149] = .22, $p = .006$)
- CISS 'compliance' problems at entry (*Spearman's rho* [149] = .20, $p = .01$)
- Shorter treatment duration (*Spearman's rho* [149] = .34, $p < .001$)
- Premature discharge (*Spearman's rho* [149] = .20, $p = .01$)

At follow-up, higher lapse frequency was associated with:

- 'Social functioning' problems (*Spearman's rho* [149] = .40, $p < .001$)
- 'Psychological functioning' problems (*Spearman's rho* [149] = .36, $p < .001$)
- 'Occupational' problems (*Spearman's rho* [149] = .20, $p = .01$)
- 'Criminal involvement' problems (*Spearman's rho* [149] = .25, $p = .002$)
- 'Drug / alcohol use' problems (*Spearman's rho* [149] = .57, $p < .001$)
- 'Ongoing support' problems (*Spearman's rho* [149] = .26, $p = .001$)
- 'Compliance' problems (*Spearman's rho* [149] = .30, $p < .001$)
- 'Working relationship' problems (*Spearman's rho* [149] = .29, $p < .001$)
- Higher CISS total scores (*Spearman's rho* [149] = .52, $p < .001$)
- 'Poor outcome' (*Spearman's rho* [149] = .43, $p < .001$)
- Lower '12 step' meeting attendance (*Spearman's rho* [149] = .28, $p = .001$)

Of the 81 patients who reported having one or more lapses, support seeking data were available for 80 of them:

- 15 patients (18.8%) sought no help for their lapse
- 22 patients (27.5%) reported using self-help groups
- 9 patients (11.3%) reported going back into Castle Craig Hospital for another period of inpatient treatment.
- 9 patients (11.3%) reported using outpatient aftercare at Castle Craig the Netherlands.
- 25 patients (31.3%) reported using some other type of support

There was no relationship between support sought and lapse frequency category within this sub-group of patients.

Detailed outcomes and what they mean for the patients

The CISS form is a rough indicator of professional impression of recent drug / alcohol related problems in the past month. Specific situations / behaviours are listed only as guiding examples and may not reflect the exact situations / behaviours of the patient. The CISS wording has been left intact in the following tables to give an idea of the actual type of dysfunction an item score of 0, 1, or 2 might indicate within each domain. The tables below illustrate the percentage of 109 patients rated as having none, moderate or severe problems within each CISS domain at intake and then again at follow-up.

Social functioning	e.g.	Intake	Follow-up
No problem	client has a stable place to live and supportive friends or relatives who are drug / alcohol free	36.5%	78.2%
Moderate problem	client's living situation may not be stable, or they may associate with drug users / heavy drinkers	43.6%	14.7%
Severe problem	living situation not stable, and they either claim to have no friends or their friends are drug users / heavy drinkers	19.9%	7.1%

General health	e.g.	Intake	Follow-up
No problem	client has reported no significant health problems	25.0%	66.0%
Moderate problem	teeth/sleep problems, occasional stomach pain, collapsed vein, asymptomatic hep B / C / HIV	57.1%	30.1%
Severe problem	extreme weight loss, jaundice, abscesses / infections, coughing up blood, fever, overdoses, blackouts, seizures, significant memory loss, neurological damage, HIV symptoms	17.9%	3.8%

Sexual or injecting risk behaviour	e.g.	Intake	Follow-up
No problem	client claims not to inject, or have unsafe sex (except in monogamous relationship with longstanding partner, spouse)	63.5%	91.7%
Moderate problem	may admit to occasional "unsafe" sexual encounters, or suspected to be injecting but denies sharing injecting equipment	29.5%	8.3%
Severe problem	client may admit to regular "unsafe" sexual encounters, or has recently been injecting and sharing injecting equipment	7.1%	0%

Psychological	e.g.	Intake	Follow-up
No problem	client appears well adjusted and relatively satisfied with the way their life is going	0%	42.9%
Moderate problem	client may have low self-esteem, general anxiety, poor sleep, may be unhappy or dissatisfied with their lot	67.9%	43.6%
Severe problem	client has a neurotic disorder e.g., panic attacks, phobias, OCD, bulimia, recently attempted or seriously considered suicide, self-harm, overdose or may be clinically depressed. Or client may have psychotic disorders, paranoia (e.g., everybody is plotting against them), deluded beliefs or hallucinations (e.g. hearing voices)	32.1%	13.5%

Occupation	e.g.	Intake	Follow-up
No problem	client is in full time occupation e.g., homemaker, parent, employed, or student	19.2%	39.7%
Moderate problem	client has some part time parenting, occupation or voluntary work	23.7%	33.3%
Severe problem	client is largely unoccupied with any socially acceptable pastime	57.1%	26.9%

Criminal involvement	e.g.	Intake	Follow-up
No problem	no criminal involvement (apart from possible possession of illicit drugs for personal use)	62.8%	93.6%
Moderate problem	client suspected of irregular criminal involvement, perhaps petty fraud, petty theft, drunk driving, small scale dealing	34.6%	5.8%
Severe problem	suspected of regular criminal involvement, or breaking and entering, car theft, robbery, violence, assault	2.6%	0.6%

Drug / alcohol use	e.g.	Intake	Follow-up
No problem	no recent drug / alcohol use	1.3%	63.5%
Moderate problem	client suspected of periodic drug / alcohol use, or else may be socially using drugs that are not considered a problem, or may be on prescribed drugs but not supplementing from other sources	9.0%	19.9%
Severe problem	client suspected of bingeing or regular drug / alcohol use	89.7%	16.7%

Ongoing support	e.g.	Intake	Follow-up
No problem	regular attendance of AA / NA, drug free drop in centre, day centre, counselling, or treatment aftercare	10.9%	43.6%
Moderate problem	patchy attendance i.e., less than once a week contact with at least one of the above	22.4%	22.4%
Severe problem	client not known to be using any type of structured support	66.7%	34.0%

Compliance	e.g.	Intake	Follow-up
No problem	attends all appointments and meetings on time, follows suggestions, or complies with treatment requirements	68.6%	82.1%
Moderate problem	not very reliable, or may have been reported as having an "attitude" problem or other difficulty with staff	24.4%	17.3%
Severe problem	chaotic, may have left treatment against staff advice or been ejected for non-compliance e.g. drug use, attitude problem	7.1%	0.6%

Working Relationship	e.g.	Intake	Follow-up
No problem	relatively easy going e.g., interviews easily, not time consuming or stressful to work with	42.9%	66.7%
Moderate problem	moderately challenging e.g., a bit demanding or time consuming, but not excessively so	53.8%	30.8%
Severe problem	quite challenging e.g., very demanding, hard work, time consuming, emotionally draining or stressful to see	3.2%	2.6%

Conclusions

- The following success rates are conservatively based by including all 201 patients on the assumption that the 45 patients not followed-up (response rate = 78%) showed no improvement or otherwise had poor outcomes.
 - Being totally abstinent from all drugs or alcohol at follow-up
49.3% ($n = 99$)
 - Achieving low problem severity at follow-up (CISS < 6, see appendix)
54.2% ($n = 109$)
 - Showing any reduction in measured levels of dysfunction
63.7% ($n = 128$)

However the success rates might be higher because some of the missing patients would not have been contactable due to having recovered and being in full time occupation.

- The following success rates are thus more liberally based by excluding the 43 patients not responding to follow-up. Although 1 patient in another addictions clinic, and 1 patient unable to answer questions because of a psychiatric condition, remained included as poor outcomes (*new sample size* = 158).
 - Being totally abstinent from all drugs or alcohol at follow-up
62.6% ($n = 99$)
 - Achieving low problem severity at follow-up (CISS < 6, see appendix)
69.0% ($n = 109$)
 - Showing any reduction in measured levels of dysfunction
81.0% ($n = 128$)

Discussion

Castle Craig Hospital provides a service to people with a range of addiction severity. Those in this group are drug or alcohol dependent people who often have the additional complications of varying degrees of co-morbidity, lack of support, poor health, and psychological problems. However, good outcomes are achieved, despite overall high levels of associated problems at intake. The patients appear similar in dysfunction to those attending outpatient drug or alcohol services. Although the goal of Castle Craig's treatment is abstinence, it should be noted that many who fail to achieve that goal still report reduced levels of problem severity at follow-up. Thus, even the treatment 'failures' appeared to have benefited from their experience in treatment, possibly by gaining a period of respite during which to recover from the consequences of their excessive drinking or drug use.

Castle Craig Hospital continues to demonstrate their ability to produce high quality research within the limitations of a busy service setting. The notion of evidence led practice is frequently discussed, but it could be argued that experienced practitioners already make best use of their resources. Thus, the purpose of such research could only be to illustrate that the experts know what they are doing (e.g., practice led evidence). This view may well be partially justified, as many of the findings in this study are obvious to those who are familiar with the field. However, some findings here are obvious only with the benefit of hindsight and others may yet inform better practice and commissioning.

References

Alford, G.S., Koehler, R.A. and Leonard, J. (1991). Alcoholics Anonymous - Narcotics Anonymous model inpatient treatment of chemically dependent adolescents: a 2-year outcome study. *Journal of Studies on Alcohol*, 52(2), 118-126.

The Audit Commission for Local Authorities and the National Health Service in England and Wales (2002). National report, changing habits: the commissioning and management of community drug treatment services for adults. Audit Commission Publications, Wetherby, UK, p71.

Christo, G. (1999a). Outcome monitoring: service evaluation made simple. GLAAS mailing, Issue 95 January 1999, Greater London Association of Alcohol Services.

Christo, G. (1999b). Keep it simple. *Drug & Alcohol Findings*, June 1999, Issue 1, p27.

Christo, G. (1999c). CISS: keeping it sweet and simple. *Addiction Today*, Vol 11, No 61, pp14-15.

Christo, G. (2000a). The Christo Inventory for Substance-misuse Services. In J. Maltby, C.A. Lewis, and A.P. Hill (Eds). *Commissioned reviews on 300 psychological tests*. Lampeter, Wales, UK: Edwin Mellen Press. p33-37.

Christo, G. (2000b). Clear classification: simple service evaluation. *Druglink (ISDD journal)*, 15(1), 19-21.

Christo, G. (2000c). CISS: keeping it sweet and simple (part 2). *Addiction Today*, Vol 11, No 62, pp14-15.

Christo, G. (2000d). Outcome monitoring must be made easy (letter to the editor). *Drug & Alcohol Findings*, issue 3, p 26. (Royal Free)

Christo, G. (2000e). Outcomes of residential and day care placements for people with drug and alcohol problems: the 2000 evaluation for Hammersmith & Fulham Social Services. Hammersmith & Fulham Social Services. (Christo Research Systems)

Christo, G. (2000f). What did they get for their money? When Hammersmith & Fulham Social Services commissioned research on addiction treatments purchased by its managers, what did it discover? Dr George Christo summarises the findings and makes recommendations to help other purchasers. *Addiction Today*, Vol. 11, No. 65, 14-15. (Christo Research Systems)

Christo, G (2001) Running numbers: dealing with the data deluge. *Druglink Vol.1*, issue 3, May/June.

Christo, G. and DaSilva, V. A. (2002). A Portuguese version of the Christo Inventory for Substance-misuse Services: a simple outcome evaluation tool. *Brazilian Journal of Medical and Biological Research*, 35, 1111-1118.

Christo, G. & Franey, C. (1995). Drug Users' Spiritual Beliefs, Locus of Control and the Disease Concept in Relation to Narcotics Anonymous Attendance and Six-Month Outcomes. *Drug and Alcohol Dependence*, 38, 51-56.

Christo, G., Spurrell, S. and Alcorn, R. (2000). Validation of the Christo Inventory for Substance-misuse Services (CISS): a simple outcome evaluation tool. *Drug and Alcohol Dependence*, 59, 189-197.

- Christo, G. and Sutton, S. (1994). Anxiety and self-esteem as a function of abstinence time among recovering addicts attending Narcotics Anonymous. *British Journal of Clinical Psychology*, 33, 198-200.
- Cook, C.H. (1988a). The Minnesota Model in the management of drug and alcohol dependency: miracle, method or myth? Part I. the philosophy and the programme. *British Journal of Addiction*, 83, 625-634.
- Cook, C.H. (1988b). The Minnesota Model in the management of drug and alcohol dependency: miracle, method or myth? Part II. evidence and conclusions. *British Journal of Addiction*, 83, 735-748.
- DeSoto, C.B., O'donnell, W.E. and DeSoto, J.L. (1989). Long-term recovery in alcoholics. *Alcoholism: Clinical and Experimental Research*, 13(5), 693-697.
- DeSoto, C.B., O'donnell, W.E., Allred, L.J. and Lopes, C.E. (1985). Symptomatology in alcoholics at various stages of abstinence. *Alcoholism: Clinical and Experimental Research*, 9(6), 505-512.
- Effective Interventions Unit (2001). Evaluation Guide 7, Using assessment data for evaluation. Effective Interventions Unit, Substance Misuse Division, Scottish Executive, St. Andrew's House, Edinburgh EH1 3DG.
- Emrick, C.D. (1987). Alcoholics Anonymous: affiliation processes and effectiveness as treatment. *Alcoholism: Clinical and Experimental Research*, 11(5), 416-423.
- Gossop M, Harris J, Best D, Man LH, Manning V, Marshall J and Strang J. (2003). Is attendance at alcoholics anonymous meetings after inpatient treatment related to improved outcomes? A 6-month follow-up study. *Alcohol Alcohol*, 38(5), 421-426.
- Longabaugh R, Wirtz PW, Zweben A, Stout RL (1998) Network support for drinking, Alcoholics Anonymous and long-term matching effects. *Addiction* 93, 1313-1333.
- Mann, R.E., Smart, R.G., Anglin, L. and Adlaf, E.M. (1991). Reductions in cirrhosis deaths in the United States: associations with per capita consumption and AA membership. *Journal of Studies on Alcohol*, 52(4), 361-365.
- McCown, W. (1989). The relationship between impulsivity, empathy and involvement in twelve step self-help substance abuse treatment groups. *British Journal of Addiction*, 84, 391-393.
- McCown, W. (1990). The effect of impulsivity and empathy on abstinence of poly- substance abusers: a prospective study. *British Journal of Addiction*, 85, 635-637.
- McLatchie, B.H. and Lomp, KGE, (1988). Alcoholics Anonymous affiliation and treatment outcome among a clinical sample of problem drinkers. *American Journal of Drug and Alcohol Abuse*, 14(3), 309-324.
- Ouimette PC, Finney JW, Moos RH (1997). Twelve-step and cognitive--behavioral treatment for substance abuse: a comparison of treatment effectiveness. *Journal of Consulting and Clinical Psychology*, 65, 230-240.
- Project MATCH (1997). Matching Alcoholism Treatments to Client Heterogeneity: Project MATCH posttreatment drinking outcomes. *Journal of Studies on Alcohol*, 58, 7-29.

Appendix, CISS comparison scores

Comparisons for interpreting CISS total score (sum of item scores)

Abstinence based treatment outcomes: Six-month outcomes for 90 treated drug users from abstinence based treatment centres

In the month before follow-up:	Good outcome: 48 were abstinent Poor outcome: 42 had used drugs	and average CISS score was 2.9 (sd = 1.9) and average CISS score was 10.6 (sd = 4.3)
Over entire six month period:	Good outcome: 33 remained abstinent* Good outcome: 22 had a lapse* Poor outcome: 35 had a relapse*	and average CISS score was 2.9 (sd = 2.0) and average CISS score was 4.5 (sd = 2.9) and average CISS score was 11.2 (sd = 4.5)

* Lapse status was assessed using an eight-level scaling of lapse / relapse outcomes (as defined by Walton et al., 1994). Drug use over the entire six-month follow-up period was assessed using the principle of Timeline Follow Back (Sobell et al., 1988), as adapted for drug use by Walton et al. (1994).

N.B. a CISS cut-off score of 6 or less can be used to indicate "good outcome" for abstinence based treatment. This correctly identified 88% of outcomes where drug use was assessed only in month before follow-up, and 84% of outcomes where drug use was assessed over the entire six-month follow-up period.

Harm minimisation prescribing based service score distribution:

Average CISS score among 243 clients at a London community drug service = 9.1 (sd = 3.4)	
16%obtained CISS scores in range 0 to 5 =	low problem severity
67%obtained CISS scores in range 6 to 12 =	average problem severity
17%obtained CISS scores in range 13 to 20=	high problem severity

Outpatient alcohol service score distribution:

Average CISS score among 102 clients at a London community alcohol service = 8.1 (sd = 3.4)	
15%obtained CISS scores in range 0 to 4 =	low problem severity
70%obtained CISS scores in range 5 to 11 =	average problem severity
15%obtained CISS scores in range 12 to 20=	high problem severity

Alcohol users generally score one CISS point less than drug users. Alcohol users are less likely to score on problems of social functioning, HIV risk behaviour and criminal involvement, but they are more likely to score on psychological problems.

References

- Christo, G. (1998). Outcomes of residential care placements for people with drug and alcohol problems. The Centre for Research on Drugs and Health Behaviour.
- Christo, G., Spurrell, S. and Alcorn, R. (2000). Validation of the Christo Inventory for Substance-misuse Services (CISS): a simple outcome evaluation tool. *Drug and Alcohol Dependence*, 59, 189-197.
- Sobell, L.C., Sobell, M.B., Leo, G.I. and Caneilla, A. (1988). Reliability of a timeline method: assessing normal drinkers' reports of recent drinking and a comparative evaluation across several populations. *British Journal of Addiction*, 83, 393-402.
- Walton, M.A., Castro, F.G. and Barrington, E.H. (1994). The role of attributions in abstinence, lapse, and relapse following substance abuse treatment. *Addictive Behaviors*, 19(3), 319-331.

CISS Website <http://users.breathemail.net/drgeorgechristo/>

© 2000 George Christo PhD, PsychD.